

AD-A087 945

IOWA UNIV IOWA CITY DEPT OF PHYSICS AND ASTRONOMY F/G 3/2
MOTION OF TRAPPED ELECTRONS AND PROTONS IN SATURN'S INNER MAGNE--ETC(U)
JUL 80 M F THOMSEN, J A VAN ALLEN N00014-76-C-0016

NL

UNCLASSIFIED

[OF]
ADA
05-79-01



END
DATE
FILMED
9-80
DTIC

JL
J. Geophys. Res., 85, in press,
1980

(12)

LEVEL

AD A087945

MOTION OF TRAPPED ELECTRONS AND PROTONS IN
SATURN'S INNER MAGNETOSPHERE

by

M. F. Thomsen and J. A. Van Allen

DTIC
ELECTE
AUG 13 1980
S D C

Department of Physics and Astronomy
The University of Iowa
Iowa City, Iowa 52242

(12) 211

July 1980

This document has been approved
for public release and sale; its
distribution is unlimited.

DDC FILE COPY

80 8 12 022

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ---	2. GOVT ACCESSION NO. AD-A087 945	3. RECIPIENT'S CATALOG NUMBER --
4. TITLE (and Subtitle) MOTION OF TRAPPED ELECTRONS AND PROTONS IN SATURN'S INNER MAGNETOSPHERE		5. TYPE OF REPORT & PERIOD COVERED Progress, July 1980
		6. PERFORMING ORG. REPORT NUMBER --
7. AUTHOR(s) M. F. Thomsen and J. A. Van Allen		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0016
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics and Astronomy The University of Iowa Iowa City, Iowa 52242		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS --
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Electronics Program Office Arlington, Virginia 22217		12. REPORT DATE July 1980
		13. NUMBER OF PAGES 20
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Submitted for publication to <u>J. Geophys. Res.</u>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Saturn's Magnetosphere Magnetospheric Physics		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) [See page following.]		

DD FORM 1473
1 JAN 73EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ABSTRACT

A summary is given of basic formulas for the guiding center motion of energetic charged particles trapped in a dipolar magnetic field. These formulas for longitudinal drift rates, latitudinal bounce periods, equatorial gyroradii, and equatorial gyroperiods are then stated in convenient numerical form for electrons and protons as functions of kinetic energy E , magnetic shell parameter L , and equatorial pitch angle α_0 for a slightly simplified model of the observed magnetic field of Saturn. To aid in the study of the interaction of charged particles with the rings and inner satellites of Saturn, additional formulas are given for the time interval between successive encounters of charged particles with a satellite in a circular prograde orbit and for the energies of electrons whose longitudinal angular velocity is resonant, or synchronous, with the Keplerian angular velocity of such a satellite.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DOC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or special
A	

INTRODUCTION

The purpose of this short paper is to summarize useful parameters which characterize the motion of charged particles in Saturn's inner magnetosphere. Such motion is a combination of gyration about a magnetic field line, latitudinal bounce along the field line, and longitudinal drift perpendicular to the field line. The longitudinal drift arises from the presence of a (corotational) electric field and from the curvature and gradient of the magnetic field. The adopted magnetic field model is that of a point dipole of moment $M = 0.20 \text{ gauss } R_s^3 = 4.32 \times 10^{28} \text{ gauss cm}^3$ (equatorial radius of the planet $= 1 R_s = 6 \times 10^9 \text{ cm}$), located at the geometrical center of the planet with \vec{M} parallel to and in the same sense as $\vec{\Omega}$, the rotational angular velocity of the planet. This model is a slight simplification of the observed field [Smith et al., 1980; Acuna and Ness, 1980]. Parameters related to the motion of electrons and protons are presented as functions of kinetic energy E , magnetic shell parameter L , and equatorial pitch angle α_0 . We adopt the convention that the longitudinal angular velocity is positive for eastward drift (same sense as the rotation of the planet) and negative for westward drift.

Unless otherwise stated, all basic formulas are in gaussian c.g.s. units and all numerical formulas give angular velocities in radian s^{-1} , distances in cm and periods in seconds when kinetic energy E is expressed in MeV. A short table of exemplary values is included. Extensive tables are contained in University of Iowa 80-25 of July 1980, available on request to the authors.

SYMBOLS AND CONSTANTS

- c = speed of light = 2.997925×10^{10} cm s^{-1}
- mc^2 = rest energy of the particle
 - = 0.511 MeV for electron.
 - = 938.3 MeV for proton
- E = kinetic energy of particle
- v = speed of particle
- β = v/c
- γ = $(1 - \beta^2)^{-\frac{1}{2}}$
- q = charge of particle = 4.80325×10^{-10} e.s.u.
per unit charge
- R = equatorial radius of planet; $1 R_s = 6 \times 10^9$ cm
for Saturn
- B_0 = equatorial surface field of planetary dipole moment
= 0.20 gauss for Saturn

M = dipole moment of planet = 4.32×10^{28} gauss cm^3
for Saturn

r = radial distance from center of the planet

α_0 = equatorial pitch angle of a particle

L = equatorial crossing distance of a dipolar
magnetic field line in units of R

λ = magnetic latitude

λ_m = magnetic latitude of a particle's mirror point

$B(\lambda), B(\lambda_m)$

= magnetic field strength at λ, λ_m

GM = gravitational constant of planet

= $3.79311 \times 10^{22} \text{ cm}^3 \text{ s}^{-2}$ for Saturn

J_2 = second zonal harmonic coefficient of planet's

gravitational potential = 1667×10^{-5} for Saturn

a = semimajor axis of a satellite orbit

Other symbols are defined as they appear in the text.

GRADIENT AND CURVATURE DRIFT

The bounce-averaged longitudinal drift of the guiding center of a particle trapped in a dipolar magnetic field due to the curvature and gradient of the field has been studied by Hamlin et al. [1961], Lew [1961] and subsequent authors. Following Hamlin et al. [1961], Lew [1961] found that the bounce-averaged angular velocity can be expressed as

$$\omega_D = \frac{3mc^3 \beta^2 \gamma L}{2qB_0 R^2} \frac{F(\lambda_m)}{G(\lambda_m)}, \quad (1)$$

where

$$F(\lambda_m) = \int_0^\pi \frac{(1 - \sin^4 \lambda)}{(1 + 3 \sin^2 \lambda)^{\frac{3}{2}}} \cdot \frac{2 - B(\lambda)/B(\lambda_m)}{[1 - B(\lambda)/B(\lambda_m)]^{\frac{1}{2}}} \cos \lambda d\lambda$$

$$G(\lambda_m) = \int_0^\pi \frac{(1 + 3 \sin^2 \lambda)^{\frac{1}{2}}}{[1 - B(\lambda)/B(\lambda_m)]^{\frac{1}{2}}} \cos \lambda d\lambda.$$

The integrals are carried out over a complete latitudinal bounce period. For Saturn, Equation (1) becomes

$$\omega_D = \pm 2.083 \times 10^{-5} LE \left(\frac{E + 2mc^2}{E + mc^2} \right) \left(\frac{F}{G} \right). \quad (2)$$

In (2) the + sign applies to protons and the - sign to electrons.

Lew [1961] further presented an analytical approximation to the ratio $F(\lambda_m)/G(\lambda_m)$ which is good to within one part in 10^3 .

$$(F/G)^{-1} = [A + B \sin^2 \lambda_m - C \exp(-k \sin^2 \lambda_m)] \quad (3)$$

with

$$A = 1.04675$$

$$B = 0.45333$$

$$C = 0.04675$$

$$k = 6.34568$$

Values of λ_m and F/G for several values of the particle's equatorial pitch angle α_0 are given in Table 1.

It is essential to note that ω_D is the longitudinal angular velocity of particle drift in the reference frame rotating with the planet.

COROTATIONAL DRIFT

In addition to longitudinal drift due to the curvature and gradient of the magnetic field, charged particles also drift in the presence of an electric field perpendicular to the magnetic field [Jackson, 1962]. Referenced to an inertial coordinate system, an electric field of this nature exists in the magnetosphere of a rotating planet if the planetary ionosphere is electrically coupled to the magnetosphere, as is the case in the presence of the ionospheric and magnetospheric plasmas which make the electrical conductivity parallel to magnetic field lines essentially infinite. Davis [1948] showed that charge flows in such a plasma surrounding a rotating, conducting magnetized sphere until the electric field

$$\vec{E} = -\frac{1}{c} (\vec{\Omega} \times \vec{r}) \times \vec{B} \quad (4)$$

is established in the plasma. This electric field results in a longitudinal drift angular velocity equal to the planet's angular velocity Ω and in the same sense [Birmingham and Northrop, 1979]. The value adopted herein for Ω , the angular velocity of Saturn's rotation, is 1.637×10^{-4} radian s^{-1} , which corresponds to the rotational period 10 hours 39.9 ± 0.3 minutes determined recently from radio observations of Saturn by Voyager instruments [Kaiser et al., 1980].

ANGULAR VELOCITY IN AN INERTIAL FRAME

The corotational angular velocity is additive to the magnetic gradient and curvature drift angular velocity. The algebraic sum of these two angular velocities is the longitudinal angular velocity of a particle relative to an inertial frame. Thus

$$\omega_I = \Omega + \omega_D . \quad (5)$$

KEPLERIAN ORBITAL ANGULAR VELOCITY

From Aksnes [1977], the mean angular velocity of a neutral body in gravitational orbit about Saturn is

$$\omega_k = \left(\frac{GM}{a^3} \right)^{\frac{1}{2}} \left(1 - \frac{3J_2 R^2}{2a^2} \right)^{-\frac{1}{2}} . \quad (6)$$

With a in units of R ,

$$\omega_k = \frac{4.1905 \times 10^{-4}}{a^{\frac{3}{2}}} \left(1 + \frac{1.25 \times 10^{-2}}{a^2} \right) . \quad (7)$$

For $r = 1.871 R_s$, $\Omega = \omega_k$; for $r > 1.871 R_s$, $\Omega > \omega_k$; and for $r < 1.871 R_s$, $\Omega < \omega_k$.

TIME INTERVAL BETWEEN SUCCESSIVE ENCOUNTERS OF A CHARGED PARTICLE WITH A SATELLITE

The angular velocity of a charged particle relative to a satellite in a circular prograde orbit in the equatorial plane at the same radial distance is

$$(\omega_I - \omega_k) = (\Omega + \omega_D - \omega_k) \quad (8)$$

and the interval between encounters is

$$T_E = 2\pi / (|\omega_I - \omega_k|) . \quad (9)$$

Note that $(\omega_I - \omega_k)$ may be positive (eastward), zero (resonant), or negative (westward).

RESONANT ENERGY

The value of E such that T_E is infinite is called the resonant (or synchronous) energy E_R and is given by

$$\Omega + \omega_D - \omega_K = 0, \quad (10)$$

a quadratic equation in E_R , whose solution for electrons is

$$E_R = 0.5 [E' - 1.022 + (E'^2 + 1.0445)^{\frac{1}{2}}], \quad (11)$$

where

$$E' = 4.800 \times 10^4 (\Omega - \omega_K) / (LF/G). \quad (12)$$

Only for $r < 1.871 R_s$ can protons be resonant with a satellite; inasmuch as this dividing radius is near the outer edge of Ring B there is no evident interest in proton resonances in Saturn's magnetosphere.

BOUNCE PERIOD

The latitudinal bounce period of a particle oscillating between mirror points on a given magnetic field line is

$$T_B = \oint \frac{ds}{v_{\parallel}}, \quad (13)$$

where the integral is carried out over arc length s along the field line for a complete (round trip) bounce between mirror points. For a dipole field,

$$T_B = \frac{4RL}{v} \int_0^{\lambda_m} \frac{\cos \lambda (1 + 3 \sin^2 \lambda)^{\frac{1}{2}}}{[1 - B(\lambda)/B(\lambda_m)]^{\frac{1}{2}}} d\lambda \quad (14)$$

$$T_B = \frac{4RL}{\beta c} H(\alpha_0) \quad (15)$$

$$T_B = 0.8006 \frac{L}{\beta} H(\alpha_0) \quad (16)$$

or

$$T_B = 0.8006 \frac{L(E + mc^2)}{[E(E + 2mc^2)]^{\frac{1}{2}}} H(\alpha_0) \quad (17)$$

The integral $H(\alpha_0)$ has been approximated by Lenchek et al. [1961]

as

$$H(\alpha_0) = 1.38 - 0.32 (\sin \alpha_0 + \sin^{\frac{1}{2}} \alpha_0) \quad (18)$$

Sample values of $H(\alpha_0)$ are given in Table 1.

EQUATORIAL GYROPERIOD

The relativistically correct angular gyrofrequency
[Jackson, 1962] of a charged particle is given by

$$\omega_g = \frac{qB}{\gamma mc} = \frac{qBc}{E + mc^2} . \quad (19)$$

In the equatorial plane of the Saturnian dipolar magnetic field,

$$\omega_g = \frac{1.798 \times 10^6}{L^3 (E + mc^2)} . \quad (20)$$

The corresponding equatorial gyroperiod $T_g = 2\pi/\omega_g$.

$$T_g = 3.495 \times 10^{-6} L^3 (E + mc^2) . \quad (21)$$

EQUATORIAL GYRORADIUS

The equatorial gyroradius (radius of the cylindrical surface on which the particle's helical trajectory lies),

$$r_g = \frac{\beta c \sin \alpha_o}{\omega_g} \quad (22)$$

or

$$r_g = 1.667 \times 10^4 L^3 \sin \alpha_o [E(E + 2mc^2)]^{\frac{1}{2}} . \quad (23)$$

APPLICABILITY OF FORMULAS

The formulas given in the foregoing sections are useful for any values of E , L , and α_0 for which the assumptions are applicable. For $L \geq 7$, the adopted model of Saturn's magnetic field becomes progressively less accurate because of magnetospheric current systems and for $L \geq 13$ the formulas are of very limited value. Also, they are not applicable to particles whose energies exceed the stable trapping limit [Van Allen, 1962].

L VALUES OF SPECIAL INTEREST

A particular purpose of this paper is to aid the study of the interaction of trapped particles with the rings and inner satellites of Saturn. L values of special interest are given in Table 2. In refined considerations, a full set of orbital elements, including eccentricity and inclination, is required [Explanatory Supplement, 1961] as are diameters of the satellites [Cruikshank, 1978].

EXEMPLARY TABLE

In Table 3, we give exemplary values of the parameters discussed herein for $L = 3.092$. Note that in this table we depart from the system of units in which the numerical formulas are given.

ACKNOWLEDGMENTS

This work was supported by the Ames Research Center/NASA contract NAS2-6553, by the U. S. Office of Naval Research, and by NASA Grant NGL 16-001-002.

Table 1

α_o	λ_m	F/G	H
90°	0.0	1.000	0.740
80	4.7	0.995	0.747
70	9.6	0.980	0.769
60	14.7	0.957	0.805
50	20.2	0.927	0.855
40	26.3	0.891	0.918
30	33.2	0.851	0.994
20	41.4	0.805	1.083
10	52.5	0.751	1.191

Table 2

L Values of Special Interest

<u>Satellite</u>	<u>L</u>	<u>References</u>
1979 S5/1979 S6	2.347	Van Allen et al. [1980b]
1979 S2/1979 S4	2.528	Van Allen et al. [1980b]
1979 S3	2.840	Van Allen et al. [1980a; 1980b]
SI Mimas	3.092	Null and Lieske [1980]; Explanatory Supplement to the Astronomical Ephemeris and Nautical Almanac [1961]
SII Enceladus	3.968	Explanatory Supplement ... [1961]
SIII Tethys	4.913	Explanatory Supplement ... [1961]
SIV Dione	6.292	Explanatory Supplement ... [1961]
SV Rhea	8.787	Explanatory Supplement ... [1961]

Table 3

Exemplary Values of Trapped Particle Parameters for $\lambda = 3.092$

All angular velocities in radian s^{-1} , T_E in hr, T_B in s, T_g in s and r_g in km.

$$\omega_k = 7.717 \text{ E} - 5, \Omega = 1.637 \text{ E} - 4$$

(a) Electrons

E (MeV)	α_0	ω_D	ω_I	$\omega_I - \omega_k$	T_E	T_B	T_g	r_g
0.1	90°	- 1.18 E - 5	1.52 E - 4	7.47 E - 5	23.4	3.34	6.31 E - 5	1.65
0.5	90	- 4.85 E - 5	1.15 E - 4	3.81 E - 5	45.9	2.12	1.04 E - 4	4.30
0.9	90	- 7.90 E - 5	8.47 E - 5	7.57 E - 6	230.	1.97	1.46 E - 4	6.48
1.005	90	- 8.65 E - 5	7.72 E - 5	Resonant	•	1.95	1.57 E - 4	7.03
1.059	60	- 8.65 E - 5	7.72 E - 5	Resonant	•	2.11	1.62 E - 4	6.34
1.218	30	- 8.65 E - 5	7.72 E - 5	Resonant	•	2.58	1.79 E - 4	4.07
1.1	90	- 9.33 E - 5	7.04 E - 5	- 6.79 E - 6	257.	1.93	1.66 E - 4	7.53
5.	90	- 3.52 E - 4	- 1.88 E - 4	- 2.65 E - 4	6.58	1.84	5.69 E - 4	27.0
10.	90	- 6.75 E - 4	- 5.12 E - 4	- 5.89 E - 4	2.96	1.83	1.09 E - 3	51.7

(b) Protons

0.1	90	1.29 E - 5	1.77 E - 4	9.94 E - 5	17.6	125.	9.70 E - 2	67.5
0.5	90	6.44 E - 5	2.28 E - 4	1.51 E - 4	11.6	56.1	9.70 E - 2	151.
1.0	90	1.29 E - 4	2.92 E - 4	2.15 E - 4	8.11	39.7	9.70 E - 2	214.
5.	90	6.42 E - 4	8.06 E - 4	7.29 E - 4	2.39	17.8	9.75 E - 2	478.
10.	90	1.28 E - 3	1.45 E - 3	1.37 E - 3	1.28	12.6	9.80 E - 2	677.
50.	90	6.28 E - 3	6.44 E - 3	6.36 E - 3	0.274	5.83	1.02 E - 1	1,530.
100.	90	1.23 E - 2	1.24 E - 2	1.23 E - 2	0.141	4.28	1.07 E - 1	2,190.

REFERENCES

- Acuna, M. H., and N. F. Ness, The magnetic field of Saturn: Pioneer 11 observations, Science, 207, 444, 1980.
- Aksnes, K., Properties of satellite orbits: Ephemerides, dynamical constants, and satellite phenomena, in Planetary Satellites, edited by J. A. Burns, University of Arizona Press, Tucson, 1977.
- Birmingham, T. J., and T. G. Northrop, Theory of flux anisotropies in a guiding center plasma, J. Geophys. Res., 84, 41, 1979.
- Cruikshank, D. P., Physical properties of the satellites of Saturn, p. 217ff of The Saturn System, NASA Conference Publication 2068, edited by D. M. Hunten and D. Morrison, 1978.
- Davis, L., Jr., Stellar electromagnetic fields, Phys. Rev., 73, 536, 1948 (abstract).
- Explanatory Supplement to the Astronomical Ephemeris and Nautical Almanac, pp. 362-386, H. M. Stationery Office, 1961.
- Hamlin, D. A., R. Karplus, R. C. Vik, and K. M. Watson, Mirror and azimuthal drift frequencies for geomagnetically trapped particles, J. Geophys. Res., 66, 1, 1961.
- Jackson, J. D., Classical Electrodynamics, John Wiley and Sons, Inc., New York, 1962.

Kaiser, M. L., M. D. Desch, J. W. Warwick, and J. B. Pearce, Voyager detection of nonthermal radio emission from Saturn, submitted to Science, 1980.

Lenchek, A. M., S. F. Singer, and R. C. Wentworth, Geomagnetically trapped electrons from cosmic ray albedo neutrons, J. Geophys. Res., 66, 4027, 1961.

Lew, J. S., Drift rate in a dipole field, J. Geophys. Res., 66, 2681, 1961.

Null, G. W., and J. H. Lieske, JPL (private communication), 1980.

Smith, E. J., L. Davis, Jr., D. E. Jones, P. J. Coleman, Jr., D. S. Colburn, P. Dyal, and C. P. Sonett, Saturn's magnetic field and magnetosphere, Science, 207, 407, 1980.

Van Allen, J. A., Dynamics, composition and origin of the geomagnetically-trapped corpuscular radiation, Trans., Int. Astronomical Union, XI B, 99, 1962.

Van Allen, J. A., B. A. Randall, and M. F. Thomsen, Sources and sinks of energetic electrons and protons in Saturn's magnetosphere, J. Geophys. Res., 85, in press, 1980a.

Van Allen, J. A., M. F. Thomsen, B. A. Randall, R. L. Rairden, and C. L. Grosskreutz, Saturn's magnetosphere, rings, and inner satellites, Science, 207, 415-421, 1980b.